

## **Hairy cell leukemia: A case report of unusual presentation without splenomegaly**

### **Shuchismita**

Department. Of Pathology (Section Hematology), Indira Gandhi Institute of Medical Sciences (IGIMS), Patna, India. Shuchi.Smita123@gmail.com

ORCID: 0000-0002-6642-6974

### **Iffat Jamal**

Department. Of Pathology (Section Hematology), Indira Gandhi Institute of Medical Sciences (IGIMS), Patna, India. Iffatjamal111@gmail.com

ORCID: 0000-0001-5574-5569

### **Vijayanand Choudhary**

Department. Of Pathology (Section Hematology), Indira Gandhi Institute of Medical Sciences (IGIMS), Patna, India. Drvijayanandchoudhary@gmail.com

ORCID: 0000-0003-1709-6128

Accepted Article

**Abstract**

Hairy cell leukemia (HCL) is characterized by pancytopenia and is usually associated with massive splenomegaly, however, the same may not be true in clinical settings. Here we report a case of HCL without the classical clinical feature of splenomegaly. Absence of splenomegaly doesn't exclude the diagnosis of HCL. A careful study of morphological findings on bone marrow aspirate and biopsy, followed by appropriate ancillary tests, aids in correct diagnosis. A high index of suspicion is essential for diagnosing and appropriately managing such cases. A strong suspicion on morphology in such cases is a prerequisite for reaching a correct diagnosis, even in the absence of usual clinical presentation.

**Keywords:** Hairy cell leukemia, Splenomegaly, Unusual, Clinical, Presentation, Bone marrow


Accepted Article

## **Introduction**

Hairy cell leukemia (HCL) is a rare B-cell neoplasm of middle age presenting with pancytopenia, splenomegaly, and hairy cells in the bone marrow. Splenomegaly accounts for 80-90% of associated features in HCL (1). A careful study of morphological findings on bone marrow aspirate and biopsy, followed by appropriate ancillary tests, aids in correct diagnosis. A high index of suspicion is essential for diagnosing and appropriately managing such cases. A strong suspicion of morphology in such cases is a prerequisite for reaching a correct diagnosis, even without the usual clinical presentation (2). Splenomegaly is by far the most constant physical finding in HCL, and HCL is typically included in the differential diagnosis of splenic enlargement. Hence a case of hairy cell leukemia presenting without splenomegaly puts both the clinician and pathologist in diagnostic dilemma many a times, as these cases show only a small percentage of abnormal mononuclear cells in the peripheral blood as well as in the bone marrow aspirate because bone marrow is usually fibrosed in these cases (3). Hence, bone marrow biopsy is the only morphologic tool available for diagnosis of HCL, highlighting the high index of suspicion that a morphologist should have in such cases, as HCL cases without splenomegaly most often have a hypocellular bone marrow, and these cases may be erroneously diagnosed as aplastic anemia (4). The incidence of HCL without splenomegaly in various series ranged from 0 to 30 % in a few of the case reports described earlier—some series report bone marrow hypocellularity to the tune of 18 % (5). Here, we represent an atypical case of HCL with weakness only but no splenomegaly. This report of an unusual presentation of such a rare disease could lead to an earlier diagnosis of HCL, even without a typical clinical presentation.

## **Case report**

A 48-year-old male presented with a three-month history of weakness and fatigue. On examination, he had pallor; however, he had no organomegaly or lymphadenopathy. He had history of on and off blood transfusion in the past. Hemogram revealed pancytopenia with hemoglobin -9.0 gm/dl, total leucocyte count- 3200/cumm with a differential count revealing 68% hairy cells which are of intermediate size and abundant blue cytoplasm, some with circumferential projections, Neutrophils- 12%, Lymphocytes -18%, Eosinophil-2%, and platelets- 90000/cumm (Figure 1a). Bone marrow aspiration (BMA) was carried out for evaluation of pancytopenia. BMA smears were hypercellular for age and showed infiltration by hairy cells. Other cell lineages were markedly suppressed (Figure 1b and c). Immunophenotyping by flow cytometry revealed a monoclonal small B-cell population with an expression of CD19 (Bright), CD20 (Bright), CD79b (Bright), CD25 (Moderate), CD103 (Moderate to bright), CD200 (dim), FMC-7(dim), CD11c (Bright), and surface kappa restricted (Moderate) (Figure 1d and e). Subsequently the patient received standard dose chemotherapy with cladribine and achieved hematological remission. Due ethical clearance was taken from institutional ethical committee with reference no IEC/IGIMS/2345. Written informed consent was taken from the patient.



**Figure 1.** a: Peripheral blood smear showing atypical lymphoid cells with circumferential projections(x100X). b: Bone marrow aspirate showing infiltration by atypical lymphoid cells. (x40x) Inset shows hypercellular marrow (x10x). c: bone marrow aspirate showing lymphoid cells with cytoplasmic projections(x100X). d and e: Flow cytometry revealing bright expression of CD11c and moderate expression of CD103 and CD25.

## Discussion

Splenomegaly is by far the most constant physical finding in HCL, and HCL is typically included in the differential diagnosis of enlarged spleen. Hence, a case of hairy cell leukemia presenting without splenomegaly puts both the clinician and pathologist in a diagnostic dilemma many times, as these cases show only a small percentage of classical abnormal mononuclear cells with circumferential projections in the peripheral blood as well as in the bone marrow aspirate (3,4).

HCL commonly presents with pancytopenia or variable cytopenia and is associated with splenomegaly and dry tap on bone marrow aspirate resulting from fibrosis (5). HCL must be included in the differential diagnoses with aplastic anemia, hypoplastic myelodysplastic syndrome, atypical chronic lymphocytic leukemia, B-prolymphocytic leukemia, and idiopathic myelofibrosis (6).

Hairy cell leukemia presenting without splenomegaly is rare. It puts both the clinician and pathologist in a diagnostic dilemma as these cases show only a small percentage of abnormal mononuclear cells in the peripheral blood as well as in the bone marrow aspirate, because bone marrow is affected by bone marrow fibrosis. Hence, bone marrow biopsy is the only morphologic tool available for diagnosis of HCL, highlighting the high index of suspicion that a morphologist should have in such cases, as HCL cases without splenomegaly most often have a hypocellular bone marrow, and these cases may be erroneously diagnosed as aplastic anemia. The incidence of HCL without splenomegaly in various series ranged from 0 to 30 %.<sup>[6,7]</sup> (Table 1).

**Table 1.** Previous studies where HCL presented without splenomegaly

Study reference	No. of patients	Absence of splenomegaly in percentage
Burke JS et al. 1974 <sup>[6]</sup>	21	0
Bouroncle et al.1979 <sup>[7]</sup>	82	7
Chatterjee et al.2008 <sup>[8]</sup>	15	0
Bhargava et al.2010 <sup>[9]</sup>	20	15
Galani et al.2012 <sup>[10]</sup>	28	4
Garman AM et al. 2013 <sup>[11]</sup>	39	30
Venkatesan et al. 2014 <sup>[12]</sup>	10	40
Present case	01	0

## **Conclusion**

This case report highlights the uncommon clinical presentation of HCL without splenomegaly creating a diagnostic challenge. A careful attention to morphological details in bone marrow biopsy is a prerequisite for its correct and early diagnosis, thereby providing maximum benefit to patients.

## **Acknowledgement**

We are grateful to the patient for providing informed consent and allowing us to share the case.

## **Funding sources**

Nil

## **Ethical statement**

This case report was approved by the institution ethics committee and all the patient information has been anonymized to protect confidentiality.

## **Conflicts of interest**

Nil

## **Author contributions**

All authors contributed to the conception and design of the case report.

## **References**

1. Zoldan MC, Ponzoni M, Ricci D, Camba L, Faravelli A. Hairy cell leukemia without splenomegaly nor myelofibrosis in a patient with gastric adenocarcinoma: early phase of the disease or a variant? *Pathologica* 87(1):97-9.
2. Venkatesan S, Purohit A, Aggarwal M, Manivannan P, Tyagi S, Mahapatra M, Pati HP, Saxena R. Unusual presentation of hairy cell leukemia: a case series of four clinically unsuspected cases. *Indian J Hematol Blood Transfus.* 2014 Sep;30(Suppl 1):413-7.
3. Muta T, Okamura T, Oshima K, Kikuchi M, Niho Y. European-American-type hairy cell leukemia without splenomegaly, treated successfully with deoxycoformycin. *Rinsho.* 2001;42(5):426-31.
4. Spedini P, Tajana M, Bergonzi C (2000) Unusual presentation of hairy cell leukemia. *Haematologica* 85(5):548-50.
5. Westbrook CA, Golde DW (1984) Clinical problems in hairy cell leukemia: diagnosis and management. *Semin Oncol* 11:514-22.
6. Burke JS, Byrne G, Appaport H (1974) Hairy cell leukemia (leukemic reticuloendotheliosis). A clinical pathologic study of 21 patients. *Cancer* 33:1394-410.
7. Bouroncle B (1979) Leukemic reticuloendotheliosis (hairy cell leukemia). *Blood* 53:412-36.
8. Chatterjee T, Panigrahi I, Mahapatra M, Pati HP, Kumar R, Naithani R et al (2008) Hairy cell leukemia: clinical, pathological and ultrastructural findings in Asian-Indians. *Indian J Cancer.* 45:41-4.
9. Bhargava R, Kar R, Mahapatra M, Saxena R (2010) The clinicohematological profile of hairy cell leukaemia: a single centre experience. *Hematology* 15:1624.
10. Galani KS, Subramanian PG, Gadage VS, Rahman K, Ashok Kumar MS, Shinde S, et. Clinico-pathological profile of Hairy cell leukemia: critical insights gained at a tertiary care cancer hospital. *Indian J Pathol Microbiol.* 2012;55:61-5.
11. Gaman AM (2013) Hairy cell leukemia—a rare type of leukemia. A retrospective study on 39 patients. *Rom J Morphol Embryol.* 54(3):575-9.

12. Venkatesan S, Purohit A, Aggrawal M, Mannivannan P, Tyagi S et al. Unusual presentation of hairy cell leukemia: A case series of four clinically unsuspected cases. *Indian J Haematol Blood Transfus.* 2014;413-7.

Accepted Article