

## The recognizing of *fli C* gene in *Pseudomonas aeruginosa* isolated from clinical sample with PCR

**Imani fooladi ,AA. (phD)**

Assistant Professor of Research  
Center of Molecular Biology,  
Baqiyatallah University of Medical  
Sciences

**Sattari ,M (phD)**

Associate Professor of Department of  
Bacteriology, Faculty of Medical  
Sciences. Tarbiat Modares University,  
Tehran, Iran

**Godarzi ,GH (MSc)**

Resident of Bacteriology, Faculty of  
Medical Sciences. Tarbiat Modares  
University, Tehran, Iran

**Corresponding Author:** Sattari M

**E.mail:** Sattarim@modares.ac.ir

### Abstract

**Background and objectives:** *Pseudomonas aeruginosa* as an opportunistic pathogen can establish lethal infections in immunocompromised patients or those exposed to predisposing factors. This bacterium contains a single polar flagellum causing motility, chemotaxis and colonization in acute phase of infection. The flagella filament is made up of a structural protein called flagellin. This study was aimed at determining The frequency of *fliC* gene in Clinical Samples.

**Material and Methods:** In this study, a pair of specific primer for types of flagellin (a, b type) was designed and by using PCR method its structural gene (*fliC*) was recognized and amplified in clinical strains.

**Results:** This original primer has appropriate efficiency in diagnostic of *pseudomonas aeruginosa* flagellum. Our study shows that 85% of the Clinical Samples have a *fliC* gene.

**Conclusion:** This method can be applied to recognizing of the motile strains, and their antigenic typing, and complete amplification of *fliC* sequence in order to cloning and expression of recombinant flagellin.

**Key words:** *Pseudomonas aeruginosa*, flagellin, *fliC*, PCR